Dr. Daniel Griffin's Clinical treatment summary for 11/16/2023

Clinical Course/Treatment – COVID

COVID PASSIVE VACCINATION

COVID POST-EXPOSURE PERIOD

https://www.cdc.gov/coronavirus/2019-ncov/your-health/quarantine-isolation.html

COVID- Early Viral - https://www.covid.gov

- 1-Paxlovid now licensed
- 2 -Remdesivir -(approved for down to 28 days of age) 3-day
- 3-Molnupiravir https://onlinelibrary.wiley.com/doi/10.1002/jmv.28011
- 4-Convalescent Plasma an early treatment option for the treatment of immunosuppressed COVID-19 patients at high risk for progression to severe disease who have no other treatment options, first week before you end up in the hospital.

ID Society on convalescent plasma "Recommending against the routine use of convalescent plasma among immunocompromised patients hospitalized with COVID-19." https://www.idsociety.org/practice-guideline/covid-19-guideline-treatment-and-management/

5-Avoid: let us not do harmful things and useless things.

6-Isolation for the Infected: Recommended 5 days.

COVID - - Cytokine Storm Week

1-Steroids at the right time in the right patient at the right dose. This is *after* the first week and in patients with oxygen saturations <94%. Dexamethasone 6mg a day x 6 days.

https://academic.oup.com/ofid/article/10/3/ofad105/7055978

- 2-Anticoagulation Guidelines from a number of organizations including ASH https://www.hematology.org/education/clinicians/guidelines-and-quality-care/clinical-practice-guidelines/venous-thromboembolism-guidelines/ash-guidelines-on-use-of-anticoagulation-in-patients-with-covid-19">https://www.hematology.org/education/clinicians/guidelines-and-quality-care/clinical-practice-guidelines/venous-thromboembolism-guidelines/ash-guidelines-on-use-of-anticoagulation-in-patients-with-covid-19
- 3-Pulmonary support.
- 4-Remdesivir if still in the first 10 days from symptom onset and not on a ventilator

Dr. Daniel Griffin's Clinical treatment summary for 11/16/2023

5-Immune modulation: Tocilizumab, the IL6-R blocker and in some cases Baricitinib, but only if there is progression and benefits outweigh risks.

6-AVOID: unnecessary antibiotics and unproven therapies